

Finding vertex-surjective graph homomorphisms*

Petr A. Golovach[†], Bernard Lidický[‡]
Barnaby Martin[§] and Daniël Paulusma[¶]

March 2, 2013

Abstract

The SURJECTIVE HOMOMORPHISM problem is to test whether a given graph G called the guest graph allows a vertex-surjective homomorphism to some other given graph H called the host graph. The bijective and injective homomorphism problems can be formulated in terms of spanning subgraphs and subgraphs, and as such their computational complexity has been extensively studied. What about the surjective variant? Because this problem is NP-complete in general, we restrict the guest and the host graph to belong to graph classes \mathcal{G} and \mathcal{H} , respectively. We determine to what extent a certain choice of \mathcal{G} and \mathcal{H} influences its computational complexity. We observe that the problem is polynomial-time solvable if \mathcal{H} is the class of paths, whereas it is NP-complete if \mathcal{G} is the class of paths. Moreover, we show that the problem is even NP-complete on many other elementary graph classes, namely linear forests, unions of complete graphs, cographs, proper interval graphs, split graphs and trees of pathwidth at most 2. In contrast, we prove that the problem is fixed-parameter tractable in k if \mathcal{G} is the class of trees and \mathcal{H} is the class of trees with at most k leaves, or if \mathcal{G} and \mathcal{H} are equal to the class of graphs with vertex cover number at most k .

1 Introduction

We consider undirected finite graphs that are *simple*, i.e., have no loops and no multiple edges. A graph is denoted $G = (V_G, E_G)$, where V_G is the set of vertices and E_G is the set of edges. A *homomorphism* from a graph G to a graph H is a mapping $f : V_G \rightarrow V_H$ that maps adjacent vertices of G to adjacent vertices of H , i.e., $f(u)f(v) \in E_H$ whenever $uv \in E_G$. Graph homomorphisms are widely studied within the areas of graph theory and algorithms; for a survey we refer to the monograph of Hell and Nešetřil [18]. The HOMOMORPHISM problem is to test whether there exists a homomorphism from a graph G called the *guest graph* to a graph H called the *host graph*. If H is restricted to be in the class of complete graphs (graphs with all possible edges), then this problem is equivalent to the COLORING problem. The latter problem is to test whether a graph G allows a k -coloring for some given k , i.e., a mapping $c : V_G \rightarrow \{1, \dots, k\}$, such that $c(u) \neq c(v)$ whenever $uv \in E_G$. This is a classical NP-complete problem [15]. Hence, the HOMOMORPHISM problem is NP-complete in general, and it is natural to restrict the input graphs to belong to some special graph classes.

We let \mathcal{G} denote the class of guest graphs and \mathcal{H} the class of host graphs that are under consideration, and denote the corresponding decision problem by $(\mathcal{G}, \mathcal{H})$ -HOMOMORPHISM. If G or H is the class of all graphs, then we use the notation “—” to indicate this. If $\mathcal{G} = \{G\}$ or $\mathcal{H} = \{H\}$, we write G and H instead of \mathcal{G} and \mathcal{H} , respectively. The Hell-Nešetřil dichotomy theorem [17] states that $(-, H)$ -HOMOMORPHISM is solvable in

*A preliminary version of this paper appeared as an extended abstract in the proceedings of CSR 2012. The work was supported by EPSRC (EP/G043434/1 and EP/G020604/1) and the Royal Society (JP090172). The 2nd author was supported by Charles University as GAUK 95710.

[†]School of Engineering and Computing Sciences, Durham University, Durham, U.K. petr.golovach@durham.ac.uk

[‡]Department of Mathematics, University of Illinois, Urbana, USA and Department of Applied Mathematics, Charles University in Prague, Czech Republic bernard@kam.mff.cuni.cz

[§]School of Engineering and Computing Sciences, Durham University, Durham, U.K. barnaby.martin@durham.ac.uk

[¶]School of Engineering and Computing Sciences, Durham University, Durham, U.K. daniel.paulusma@durham.ac.uk

polynomial time if H is bipartite, and NP-complete otherwise. In the context of graph homomorphisms, a graph F is called a *core* if there exists no homomorphism from F to any proper subgraph of F . Dalmau et al. [5] proved that the $(\mathcal{G}, -)$ -HOMOMORPHISM problem can be solved in polynomial time if all cores of the graphs in \mathcal{G} have bounded treewidth. Moreover, Grohe [16] showed that under the assumption $\text{FPT} \neq \text{W}[1]$, the problem can be solved in polynomial time if and only if this condition holds.

As a homomorphism f from a graph G to a graph H is a (vertex) mapping, we may add further restrictions, such as requiring it to be *bijective*, *injective*, or *surjective* i.e., for each $x \in V_H$ there exists exactly one, at most one, or at least one vertex $u \in V_G$ with $f(u) = x$, respectively. The decision problems corresponding to the first and second variant are known as the SPANNING SUBGRAPH ISOMORPHISM and SUBGRAPH ISOMORPHISM problem, respectively. As such, these two variants have been well studied in the literature. For example, the bijective variant contains the problem that is to test whether a graph contains a Hamiltonian cycle as a special case. In our paper, we research the third variant, which leads to the following decision problem:

SURJECTIVE HOMOMORPHISM

Instance: two graphs G and H .

Question: does there exist a surjective homomorphism from G to H ?

If the guest G is restricted to a graph class \mathcal{G} and the host H to a graph class \mathcal{H} , then we denote this problem by SURJECTIVE $(\mathcal{G}, \mathcal{H})$ -HOMOMORPHISM. Fixing the host side to a single graph H yields the SURJECTIVE $(-, H)$ -HOMOMORPHISM problem. This problem is NP-complete already when H is nonbipartite. This follows from a simple reduction from the corresponding $(-, H)$ -HOMOMORPHISM problem, which is NP-complete due to the Hell-Nešetřil dichotomy theorem [17]; we replace an instance graph G of the latter problem by the disjoint union $G + H$ of G and H , and observe that G allows an homomorphism to H if and only if $G + H$ allows a surjective homomorphism to H . For bipartite host graphs H , the complexity classification of SURJECTIVE $(-, H)$ -HOMOMORPHISM is still open, although some partial results are known. For instance, the problem can be solved in polynomial time whenever H is a tree. This follows from a more general classification that also includes trees in which the vertices may have self-loops [14]. On the other hand, there exist cases of bipartite host graphs H for which the problem is NP-complete, e.g., when H is the graph obtained from a 6-vertex cycle with one distinct path of length 3 added to each of its six vertices [2]. Recently, the SURJECTIVE $(-, H)$ -HOMOMORPHISM problem has been shown to be NP-complete when H is a 4-vertex cycle with a self-loop in every vertex [20]. Note that in our paper we only consider simple graphs. For a survey on the SURJECTIVE $(-, H)$ -HOMOMORPHISM problem from a constraint satisfaction point of view we refer to the paper of Bodirsky, Kára and Martin [2]. Below we discuss some other concepts that are closely related to surjective homomorphisms.

A homomorphism f from a graph G to a graph H is *locally surjective* if f becomes surjective when restricted to the neighborhood of every vertex u of G , i.e., $f(N_G(u)) = N_H(f(u))$. The corresponding decision is called the ROLE ASSIGNMENT problem which has been classified for any fixed host H [11]. Any locally surjective homomorphism is surjective if the host graph is connected but the reverse implication is not true in general. For more on locally surjective homomorphisms and the locally injective and bijective variants, we refer to the survey of Fiala and Kratochvíl [9].

Let H be an induced subgraph of a graph G . Then a homomorphism f from a graph G to H is a *retraction* from G to H if $f(h) = h$ for all $h \in V_H$. In that case we say that G *retracts to* H . By definition, a retraction from G to H is a surjective homomorphism from G to H . Retractions are well studied; see e.g. the recent complexity classification of Feder et al. [7] for the corresponding decision problem when H is a fixed pseudoforest. In particular, polynomial-time algorithms for retractions have been proven to be a useful subroutine for obtaining polynomial-time algorithms for the SURJECTIVE $(-, H)$ -HOMOMORPHISM problem [14].

We emphasize that a surjective homomorphism is *vertex-surjective* as opposed to the stronger condition of being *edge-surjective*. A homomorphism from a graph G to a graph H is called *edge-surjective* or a *compaction* if for any edge $xy \in E_H$ there exists an edge $uv \in E_G$ with $f(u) = x$ and $f(v) = y$. If f is a compaction from G to H , we also say that G *compacts to* H . The COMPACTION problem is to test whether a graph G compacts to a graph H . Vikas [21, 22, 23] determined the computational complexity

\mathcal{G}	\mathcal{H}	Complexity	
complete graphs	all graphs	polynomial time	Proposition 1 (i)
all graphs	paths	polynomial time	Proposition 1 (ii)
paths	all graphs	NP-complete	Theorem 1 (i)
linear forests	linear forests	NP-complete	Theorem 1 (ii)
unions of complete graphs	unions of complete graphs	NP-complete	Theorem 1 (iii)
connected cographs	connected cographs	NP-complete	Theorem 1 (iv)
trees of $\mathbf{pw} \leq 2$	trees of $\mathbf{pw} \leq 2$	NP-complete	Theorem 1 (v)
split graphs	split graphs	NP-complete	Theorem 1 (vi)
connected proper interval graphs	connected proper interval graphs	NP-complete	Theorem 1 (vii)
trees	trees with k leaves	FPT in k	Theorem 2
graphs of $\mathbf{vc} \leq k$	graphs of $\mathbf{vc} \leq k$	FPT in k	Theorem 3

Table 1: Complexity of $(\mathcal{G}, \mathcal{H})$ -SURJECTIVE HOMOMORPHISM.

of $(-, H)$ -COMPACTION for several classes of fixed host graphs H . Very recently, Vikas [24] considered $(-, H)$ -COMPACTION for guest graphs belonging to some restricted graph class.

Our Results. We study the SURJECTIVE $(\mathcal{G}, \mathcal{H})$ -HOMOMORPHISM problem for several graph classes \mathcal{G} and \mathcal{H} . We observe that this problem is polynomial-time solvable when the host graph is a path, whereas it becomes NP-complete if we restrict the guests to be paths instead of the hosts. We also show that the problem is NP-complete when both \mathcal{G} and \mathcal{H} are restricted to trees of pathwidth at most 2, and when both \mathcal{G} and \mathcal{H} are linear forests. These results are in contrast to the aforementioned polynomial-time result of Dalmau et al. [5] on $(\mathcal{G}, -)$ -HOMOMORPHISM for graph classes \mathcal{G} that consists of graphs, the cores of which have bounded treewidth. They are also in contrast to the aforementioned polynomial-time result on SURJECTIVE $(-, H)$ -HOMOMORPHISM when H is any fixed tree [14].

Due to the hardness for graphs of bounded treewidth, it is natural to consider other width parameters such as the clique-width of a graph. For this purpose we first consider the class of complete graphs that are exactly those graphs that have clique-width 1. We observe that the SURJECTIVE $(\mathcal{G}, \mathcal{H})$ -HOMOMORPHISM can be solved in polynomial time when \mathcal{G} is the class of complete graphs, whereas the problem becomes NP-complete when we let \mathcal{G} and \mathcal{H} consist of the unions of complete graphs. We then focus on graphs that have clique-width at most two. This graph class is equal to the class of cographs [4]. There exist only a few natural problems that are difficult on cographs. We prove that SURJECTIVE $(\mathcal{G}, \mathcal{H})$ -HOMOMORPHISM, where \mathcal{G} and \mathcal{H} are equal to the class of connected cographs, is one of these. We also consider proper interval graphs. This graph class has unbounded tree-width and contains the classes of complete graphs and paths. Because they are “path-like”, often problems that are difficult for general graphs are tractable for proper interval graphs. In an attempt to generalize our polynomial-time result for SURJECTIVE $(\mathcal{G}, \mathcal{H})$ -HOMOMORPHISM when \mathcal{G} is the class of complete graphs, or when \mathcal{H} is the class of paths, we consider connected proper interval graphs. It turns out that SURJECTIVE $(\mathcal{G}, \mathcal{H})$ -HOMOMORPHISM is NP-complete even when \mathcal{G} and \mathcal{H} consist of these graphs. Our last hardness result shows that the problem is also NP-complete when \mathcal{G} and \mathcal{H} are equal to the class of split graphs. All hardness results can be found in Section 3.

To complement our hardness results, we show in Section 4 that SURJECTIVE $(\mathcal{G}, \mathcal{H})$ -HOMOMORPHISM is fixed-parameter tractable in k , when \mathcal{G} is the class of trees and \mathcal{H} is the class of trees with at most k leaves, and also when \mathcal{G} and \mathcal{H} consist of graphs with vertex cover number at most k . The latter result adds further evidence that decision problems difficult for graphs of bounded treewidth may well be tractable if the vertex cover number is bounded; also see e.g. [1, 6, 8, 10]. Moreover, the vertices of such graphs can be partitioned into two sets, one of them has size bounded by the vertex cover number and the other one is an independent set. As such, they resemble split graphs with bounded clique number. We refer to Table 1 for a summary of our results. In this table, \mathbf{pw} and \mathbf{vc} denote the pathwidth and the vertex cover number of a graph, respectively. In Section 2 we explain these notions and the complexity class FPT. There, we also give the definitions of all the aforementioned graph classes.

2 Definitions and Preliminaries

Let G be a graph. The *open neighborhood* of a vertex $u \in V_G$ is defined as $N_G(u) = \{v \mid uv \in E_G\}$, and its *closed neighborhood* is defined as $N_G[u] = N_G(u) \cup \{u\}$. The degree of a vertex $u \in V_G$ is denoted $d_G(u) = |N_G(u)|$. The *distance* $\text{dist}_G(u, v)$ between a pair of vertices u and v of G is the number of edges of a shortest path between them. The *distance* between a vertex u and a set of vertices $S \subseteq V_G$ is $\text{dist}_G(u, S) = \min\{\text{dist}_G(u, v) \mid v \in S\}$. We may omit subscripts if this does not create any confusion. The *diameter* of G is defined as $\text{diam}(G) = \max\{\text{dist}_G(u, v) \mid u, v \in V_G\}$. Let $S \subseteq V_G$. Then the graph $G - S$ is the graph obtained from G by removing all vertices in S . If $S = \{u\}$, we also write $G - u$. The subgraph of G that is *induced* by S has vertex set S and edges uv if and only if $uv \in E_G$. We denote this subgraph by $G[S]$.

A graph is an *interval graph* if intervals of the real line can be associated with its vertices in such a way that two vertices are adjacent if and only if their corresponding intervals overlap. An interval graph is *proper* if it has an interval representation, in which no interval is properly contained in any other interval. The disjoint union of two graphs G and H is denoted $G + H$, and the disjoint union of r copies of G is denoted rG . A *linear forest* is the disjoint union of a collection of paths. We denote the path on n vertices by P_n . A graph is a *cograph* if it does not contain P_4 as an induced subgraph. A *clique* is the vertex set of a complete graph. A vertex set is *independent* if its vertices are mutually non-adjacent. A graph is a *split graph* if its vertex set can be partitioned into a clique and an independent set.

A *tree decomposition* of a graph G is a pair (\mathcal{X}, T) where T is a tree and $\mathcal{X} = \{X_i \mid i \in V_T\}$ is a collection of subsets (called *bags*) of V_G such that the following three conditions are satisfied:

1. $\bigcup_{i \in V_T} X_i = V_G$;
2. for each edge $xy \in E_G$, the vertices x, y are in a bag X_i for some $i \in V_T$;
3. for each $x \in V_G$, the set $\{i \mid x \in X_i\}$ induces a connected subtree of T .

The *width* of tree decomposition (\mathcal{X}, T) is $\max_{i \in V_T} \{|X_i| - 1\}$. The *treewidth* of a graph G , denoted $\text{tw}(G)$, is the minimum width over all tree decompositions of G . If in these two definitions we restrict the tree T to be a path, then we obtain the notions of *path decomposition* and *pathwidth* of G denoted $\text{pw}(G)$.

For a graph G , a set $S \subseteq V_G$ is a *vertex cover* of G , if every edge of G has at least one of its two endvertices in S . Let $\text{vc}(G)$ denote the *vertex cover number*, i.e., the minimum size of a vertex cover of G .

We use the following well-known notion in parameterized complexity, where one considers the problem input as a pair (I, k) , where I is the main part and k the parameter; also see the text book of Flum and Grohe [12]. A problem is *fixed parameter tractable* if an instance (I, k) can be solved in time $O(f(k)n^c)$, where f denotes a computable function, n denotes the size of I , and c is a constant independent of k . The class FPT is the class of all fixed-parameter tractable decision problems.

We finish this section by giving the polynomial-time results from Table 1. The proof of statement (ii) of Proposition 1 is similar to the corresponding proof for the edge-surjective variant shown by Vikas [24].

Proposition 1. *The SURJECTIVE $(\mathcal{G}, \mathcal{H})$ -HOMOMORPHISM problem can be solved in polynomial time in the following two cases:*

- (i) \mathcal{G} is the class of complete graphs and \mathcal{H} is the class of all graphs;
- (ii) \mathcal{G} is the class of all graphs and \mathcal{H} is the class of paths.

Proof. We first prove (i). Let G be a complete graph and H be an arbitrary graph. We claim that there exists a surjective homomorphism from G to H if and only if H is a complete graph with the same number of vertices as G . Because this condition can be checked in polynomial time, showing this is sufficient to prove (i).

First suppose that H is a complete graph with the same number of vertices as G . Then the identity mapping is a surjective homomorphism from G to H .

Now suppose that f is a surjective homomorphism from G to H . Because G is a complete graph and f is a homomorphism, there are no two distinct vertices u and v with $f(u) = f(v)$. Because f is surjective, this means that $|V_G| = |V_H|$. Let x and y be two distinct vertices in H . Because f is surjective, there exist two vertices u and v in G with $f(u) = x$ and $f(v) = y$. Because G is a complete graph, u and v are adjacent. Then, because f is a homomorphism, x and y must be adjacent. Hence, H is a complete graph. This completes the proof of (i).

We now prove (ii). Suppose that we are given a guest graph G with k connected components G_1, \dots, G_k for some $k \geq 1$, and a host path P_ℓ for some $\ell \geq 1$. If $\ell = 1$, then there exists a surjective homomorphism from G to P_ℓ if and only if each G_i consists of one vertex. Assume that $\ell \geq 2$. We claim that there exists a surjective homomorphism from G to P_ℓ if and only if a) G is bipartite and b) $\sum_{i=1}^k \text{diam}(G_i) + k \geq \ell$. Because conditions a) and b) can be checked in polynomial time, showing this is sufficient to prove (ii).

First suppose that f is a surjective homomorphism from G to P_ℓ . Because P_ℓ is a bipartite graph, G is bipartite as well, and a) holds. For $i = 1, \dots, k$, we let P^i denote the subgraph of P_ℓ induced by $f(V_{G_i})$. Because each G_i is connected and f is a homomorphism, each P^i is connected, and hence, forms a subpath of P_ℓ . Because f is a homomorphism, $\text{diam}(G_i) \geq |V_{P^i}| - 1$ for $i = 1, \dots, k$. We use this inequality and the surjectivity of f to obtain

$$\ell \leq \sum_{i=1}^k |V_{P^i}| \leq \sum_{i=1}^k (\text{diam}(G_i) + 1) = \sum_{i=1}^k \text{diam}(G_i) + k.$$

Now suppose that G is bipartite and that $\sum_{i=1}^k \text{diam}(G_i) + k \geq \ell$. Let $F = G_i$ be an arbitrary connected component of G . We first prove that for all $\min\{1, \text{diam}(F)\} + 1 \leq s \leq \text{diam}(F) + 1$, there is a surjective homomorphism h from F to P_s . Clearly, this holds if $\text{diam}(F) = 0$. Let $\text{diam}(F) \geq 1$. Then $s \geq 2$. Let $P = v_1 v_2 \dots v_s$. Let u be a vertex of F such that F has a vertex at distance $\text{diam}(F)$ from u . We consider a mapping $h: V_F \rightarrow \{v_1, \dots, v_k\}$ such that $h(x) = v_i$, where

$$i = \begin{cases} \text{dist}_F(u, x) + 1 & \text{if } \text{dist}_F(u, x) \leq s - 2, \\ s & \text{if } \text{dist}_F(u, x) \geq s - 1 \text{ and } (\text{dist}_F(u, x) - s + 1) \bmod 2 = 0, \\ s - 1 & \text{if } \text{dist}_F(u, x) \geq s - 1 \text{ and } (\text{dist}_F(u, x) - s + 1) \bmod 2 = 1. \end{cases}$$

Because G is bipartite, F is bipartite. Then h is a homomorphism, and because $\text{diam}(G) \geq s - 1$, h is surjective.

Because $\sum_{i=1}^k \text{diam}(G_i) + k \geq \ell \geq 2$, we can cover P_ℓ by subpaths P^1, \dots, P^k (i.e., $\cup_{i=1}^k V_{P^i} = V_{P_\ell}$) in such a way that for all $1 \leq i \leq k$, we have that $\min\{1, \text{diam}(G_i)\} + 1 \leq |V_{P^i}| \leq \text{diam}(G_i) + 1$. It remains to recall that we have a surjective homomorphism from each G_i to P^i , and claim b) follows. This completes the proof of (ii), and hence, we have shown Proposition 1. \square \square

3 Hard Cases

In contrast to case (ii) of Proposition 1, where the host graphs are assumed to be paths, our problem becomes difficult when the guest graphs are restricted to paths. Our next theorem shows this and the other hardness results of Table 1.

Theorem 1. *The SURJECTIVE $(\mathcal{G}, \mathcal{H})$ -HOMOMORPHISM problem is NP-complete in the following six cases:*

- (i) \mathcal{G} is the class of paths and \mathcal{H} is the class of all graphs;
- (ii) $\mathcal{G} = \mathcal{H}$ is the class of linear forests;
- (iii) $\mathcal{G} = \mathcal{H}$ is the class of disjoint unions of complete graphs;
- (iv) $\mathcal{G} = \mathcal{H}$ is the class of connected cographs;




Figure 1: The linear forests G and H constructed in the proof of (ii).

(v) $\mathcal{G} = \mathcal{H}$ is the class of trees of pathwidth at most two;

(vi) $\mathcal{G} = \mathcal{H}$ is the class of split graphs;

(vii) $\mathcal{G} = \mathcal{H}$ is the class of connected proper interval graphs.

Proof. We first prove (i). We reduce from the well-known problem HAMILTONIAN PATH, which is NP-complete [15]. An n -vertex graph H has a Hamiltonian path if and only if there exists a surjective homomorphism from P_n to H . This proves (i).

For showing (ii)-(vii) we need some extra terminology. We say that a multiset $A = \{a_1, \dots, a_n\}$ of integers is (m, B) -positive if $n = 3m$, $\sum_{i=1}^n a_i = mB$ and $a_i > 0$ for $i = 1, \dots, n$. A 3-partition of a multiset $A = \{a_1, \dots, a_n\}$ that is (m, B) -positive for some integers m, B is a partition S_1, S_2, \dots, S_m of A such that for $1 \leq j \leq m$, $|S_j| = 3$ and $\sum_{a_i \in S_j} a_i = B$. This leads to the problem:

3-PARTITION

Instance: an (m, B) -positive multiset $A = \{a_1, \dots, a_n\}$ for some integers m, B ;

Question: does A have a 3-partition?

The 3-PARTITION problem is known to be NP-complete [15] in the strong sense, i.e., it remains hard even if all integers in the input are encoded in unary. This enables us to reduce from this problem in order to show NP-completeness in the cases (ii)-(vii). In each of these six cases we assume that $A = \{a_1, \dots, a_n\}$ is a (m, B) -positive multiset for some integers m, B . We now prove (ii)-(vii).

(ii) For $i = 1, \dots, n$, let $p_i = a_i + B$, and let $q = 4B$. Let G be the linear forest $G_1 + \dots + G_n$, where G_i is isomorphic to P_{p_i} for $i = 1, \dots, n$. Let H be the linear forest $H_1 + \dots + H_m = mP_q$. The forests G and H are displayed in Figure 1. We claim that A has a 3-partition if and only if there exists a surjective homomorphism from G to H .

Suppose that S_1, \dots, S_m is a 3-partition of A . For each $1 \leq j \leq m$, we consider the connected components $G_{i_1}, G_{i_2}, G_{i_3}$ of G such that $S_j = \{a_{i_1}, a_{i_2}, a_{i_3}\}$. We map the vertices of G_{i_1} to the first p_{i_1} vertices of H_j according to the path order, and similarly the vertices of G_{i_2} to the next p_{i_2} vertices of H_j , and the vertices of G_{i_3} to the last p_{i_3} vertices of H_j . Because $p_{i_1} + p_{i_2} + p_{i_3} = a_{i_1} + a_{i_2} + a_{i_3} + 3B = 4B = q$ for $i = 1, \dots, n$, we obtain a surjective homomorphism from G to H in this way.

Now suppose that f is a surjective homomorphism from G to H . We observe that $|V_G| = |V_H| = 4mB$. Hence, f is also injective. Because f is a homomorphism, f must map all vertices of each connected component of G to the same connected component of H . Let $1 \leq j \leq m$, and let G_{i_1}, \dots, G_{i_s} be the connected components of G that are mapped to H_j . Because $|V_{H_j}| = 4B$ and every connected component of G contains at least $B + 1$ vertices, we find that $s \leq 3$. Because G has $3m$ connected components, we then find that $s = 3$. Because f is injective, $a_{i_1} + a_{i_2} + a_{i_3} + 3B = p_{i_1} + p_{i_2} + p_{i_3} = q = 4B$. Hence, $a_{i_1} + a_{i_2} + a_{i_3} = B$ and we let $S_j = \{a_{i_1}, a_{i_2}, a_{i_3}\}$. This means that we obtain the partition S_1, \dots, S_m of A that is a 3-partition. This completes the proof of (ii).

(iii) We use all arguments from the proof of (ii) after replacing each path in G and H by a clique of the same size.

(iv) In the graphs G and H from the proof of (ii) we replace each path by a clique of the same size. We also add a vertex v in G adjacent to all other vertices of G , and a vertex x in H adjacent to all other vertices of




Figure 2: The trees G and H constructed in the proof of (v).

H . The resulting graphs are connected cographs. We observe that every homomorphism maps v to x . To finish the proof we use the same arguments as the ones used to prove (ii).

(v) For $i = 1, \dots, n$, let $p_i = a_i + B$, and let $q = 4B$. We construct two trees G and H . We first construct G :

- for $i = 1, \dots, n$, introduce p_i vertices $u_1^{(i)}, \dots, u_{p_i}^{(i)}$ and a vertex v_i adjacent to $u_1^{(i)}, \dots, u_{p_i}^{(i)}$;
- add a new vertex w and make it adjacent to v_1, \dots, v_n .

We now construct H :

- for $j = 1, \dots, m$, introduce q vertices $x_1^{(j)}, \dots, x_q^{(j)}$ and a vertex y_j adjacent to $x_1^{(j)}, \dots, x_q^{(j)}$;
- add a new vertex z and make it adjacent to y_1, \dots, y_m .

The trees G and H are displayed in Figure 2. For G we take the path decomposition with bags $\{u_h^{(i)}, v_i, w\}$ to find that $\mathbf{pw}(G) \leq 2$. Similarly, we find that $\mathbf{pw}(H) \leq 2$. We claim that A has a 3-partition if and only if there is a surjective homomorphism from G to H .

First suppose that S_1, \dots, S_m is a 3-partition of A . We define f as follows. We set $f(w) = z$. Then for $j = 1, \dots, m$, we consider the set $S_j = \{a_{i_1}, a_{i_2}, a_{i_3}\}$. We let f map the vertices $v_{i_1}, v_{i_2}, v_{i_3}$ to y_j . Then we let f map the vertices $u_1^{(i_1)}, \dots, u_{p_{i_1}}^{(i_1)}$ consecutively to the first p_{i_1} vertices of the set $\{x_1^{(j)}, \dots, x_q^{(j)}\}$, the vertices $u_1^{(i_2)}, \dots, u_{p_{i_2}}^{(i_2)}$ to the next p_{i_2} vertices of this set, and finally, the vertices $u_1^{(i_3)}, \dots, u_{p_{i_3}}^{(i_3)}$ to the last p_{i_3} vertices of the set. Because $p_{i_1} + p_{i_2} + p_{i_3} = a_{i_1} + a_{i_2} + a_{i_3} + 3B = 4B = q$, we find that f is a surjective homomorphism from G to H .

Now suppose that f is a surjective homomorphism from G to H . We observe that $f(w) = z$, because all vertices of G must be mapped at distance at most two from $f(w)$. Consequently, f maps every v -vertex to a y -vertex, and every u -vertex to an x -vertex. The number of u -vertices is $p_1 + \dots + p_n = a_1 + \dots + a_n + nB = 4mB$, which is equal to the number of x -vertices. Hence f maps the u -vertices bijectively to the x -vertices. Moreover, if $f(v_i) = y_j$, then f maps the vertices $u_1^{(i)}, \dots, u_{p_i}^{(i)}$ to the vertices from the set $\{x_1^{(j)}, \dots, x_q^{(j)}\}$. For $j = 1, \dots, m$, let v_{i_1}, \dots, v_{i_s} be the vertices mapped to y_j . Because $p_i > B$ for all $1 \leq i \leq n$, we find that $s \leq 3$. Then, because $n = 3m$, we conclude that $s = 3$. Because f maps bijectively $\{u_1^{(i_1)}, \dots, u_{p_{i_1}}^{(i_1)}\} \cup \{u_1^{(i_2)}, \dots, u_{p_{i_2}}^{(i_2)}\} \cup \{u_1^{(i_3)}, \dots, u_{p_{i_3}}^{(i_3)}\}$ to $\{x_1^{(j)}, \dots, x_q^{(j)}\}$, we find that $a_{i_1} + a_{i_2} + a_{i_3} + 3B = p_{i_1} + p_{i_2} + p_{i_3} = q = 4B$, and consequently, $a_{i_1} + a_{i_2} + a_{i_3} = B$. We set $S_j = \{a_{i_1}, a_{i_2}, a_{i_3}\}$. It remains to observe that S_1, \dots, S_m is a 3-partition of A . This completes the proof of (v).

(vi) For $i = 1, \dots, n$, let $p_i = a_i + B$, and let $q = 4B$. We construct two graphs G and H . We first construct G :

- for $i = 1, \dots, n$, introduce p_i vertices $u_1^{(i)}, \dots, u_{p_i}^{(i)}$ and a vertex v_i adjacent to $u_1^{(i)}, \dots, u_{p_i}^{(i)}$;




Figure 3: The split graphs G and H constructed in the proof of (vi).




Figure 4: The proper interval graphs G and H constructed in the proof of (vii).

- joint all u -vertices by edges pairwise to obtain a clique of size $4mB$.

We construct H as follows:

- for $j = 1, \dots, m$, introduce q vertices $x_1^{(j)}, \dots, x_q^{(j)}$ and vertices $y_j^{(1)}, y_j^{(2)}, y_j^{(3)}$ adjacent to $x_1^{(j)}, \dots, x_q^{(j)}$;
- joint all x -vertices by edges pairwise to obtain a clique of size $4mB$.

We observe that G and H are split graphs, also see Figure 3. We claim that A has a 3-partition if and only if there is a surjective homomorphism from G to H .

First suppose that S_1, \dots, S_m is a 3-partition of A . We define f as follows. For $j = 1, \dots, m$, we consider the set $S_j = \{a_{i_1}, a_{i_2}, a_{i_3}\}$. We let f map the vertices $v_{i_1}, v_{i_2}, v_{i_3}$ to $y_j^{(1)}, y_j^{(2)}, y_j^{(3)}$ respectively. Then we let f map the vertices $u_1^{(i_1)}, \dots, u_{p_{i_1}}^{(i_1)}$ to the first p_{i_1} vertices of the set $\{x_1^{(j)}, \dots, x_q^{(j)}\}$, the vertices $u_1^{(i_2)}, \dots, u_{p_{i_2}}^{(i_2)}$ to the next p_{i_2} vertices of this set, and finally, the vertices $u_1^{(i_3)}, \dots, u_{p_{i_3}}^{(i_3)}$ to the last p_{i_3} vertices of the set. Because $p_{i_1} + p_{i_2} + p_{i_3} = a_{i_1} + a_{i_2} + a_{i_3} + 3B = 4B = q$, we find that f is a surjective homomorphism from G to H .

Now suppose that f is a surjective homomorphism from G to H . Observe that $|V_G| = |V_H|$. Hence, f is a bijection. The homomorphism f maps any clique of G to a clique of the same size in H . It follows that all u -vertices of G are mapped to x -vertices of H , and all v -vertices of G are mapped to y -vertices of H . For $j = 1, \dots, m$, let $v_{i_1}, v_{i_2}, v_{i_3}$ be the vertices mapped to $y_j^{(1)}, y_j^{(2)}, y_j^{(3)}$ respectively. Then the vertices $u_1^{(i_1)}, \dots, u_{p_{i_1}}^{(i_1)}$, $u_1^{(i_2)}, \dots, u_{p_{i_2}}^{(i_2)}$ and $u_1^{(i_3)}, \dots, u_{p_{i_3}}^{(i_3)}$ are mapped bijectively to the vertices $x_1^{(j)}, \dots, x_q^{(j)}$. Therefore, $a_1 + a_2 + a_3 + 3B = p_1 + p_2 + p_3 = q = 4B$ and $a_1 + a_2 + a_3 = B$. We set $S_j = \{a_{i_1}, a_{i_2}, a_{i_3}\}$, and it remains to observe that S_1, \dots, S_m is a 3-partition of A . This completes the proof of (vi).

(vii) For $i = 1, \dots, n$, let $p_i = 6m^2(a_i + B)$, and let $q = 24m^2B$. We construct two graphs G and H . We first construct G :

- for $i = 1, \dots, n$, construct a clique U_i on p_i vertices $u_1^{(i)}, \dots, u_{p_i}^{(i)}$;
- for $i = 2, \dots, n$, join $u_{p_{i-1}}^{(i-1)}$ and $u_1^{(i)}$ by a path Q_i of length $2m - 1$.

Then we construct H :

- for $j = 1, \dots, m$, construct a clique $X^{(j)}$ on q vertices $x_1^{(j)}, \dots, x_q^{(j)}$;
- for $j = 2, \dots, m$, join $x_q^{(j-1)}$ and $x_1^{(j)}$ by an edge.

We observe that G and H are proper interval graphs, also see Figure 4. We claim that A has a 3-partition if and only if there exists a surjective homomorphism from G to H .

First suppose that S_1, \dots, S_m is a 3-partition of A . We partition each $X^{(j)}$ into three cliques $X_{i_1} \cup X_{i_2} \cup X_{i_3}$ of size p_{i_1} , p_{i_2} , and p_{i_3} , respectively, corresponding to $S_j = \{a_{i_1}, a_{i_2}, a_{i_3}\}$; this is possible because $|X^{(j)}| = q = 24m^2B = 6m^2(a_{i_1} + a_{i_2} + a_{i_3} + 3B) = p_{i_1} + p_{i_2} + p_{i_3}$; We will determine a homomorphism f from G to H such that f is a bijection from U_i to X_i for $i = 1, \dots, n$. Hence, this property will ensure that f is surjective. In order to do this, we must show that we do not violate the definition of a homomorphism with respect to the remaining vertices of G ; note these remaining vertices are the inner vertices of the Q -paths. We therefore define f inductively as follows.

Let $i = 1$. Assume that $a_1 \in S_j$. We let f map the vertices of U_1 to the vertices of X_1 bijectively in an arbitrary order.

Let $i \geq 2$ and suppose that f is constructed for all vertices of U_s and Q_s for all $1 \leq s \leq i-1$. Let $y = f(u_{p_{i-1}}^{(i-1)})$. Because H has diameter at most $2m-1$, we find that y is at distance at most $2m-1$ from the set X_i . Consider the subgraph H' of H that contains X_i and a shortest path between y and X_i . Because $|X_i| \geq 2m$, we find that H' contains a (y, z) -path of length $2m-1$ for some vertex $z \in X_i$. Recall that $y = f(u_{p_{i-1}}^{(i-1)})$. We map consecutively the vertices of the $(u_{p_{i-1}}^{(i-1)}, u_1^{(i)})$ -path Q_i of length $2m-1$ to the vertices of P in the path order. Note that $f(u_1^{(i)}) = z$. Then we map the vertices $u_2^{(i)}, \dots, u_{p_i}^{(i)}$ to the vertices of $X_i \setminus \{z\}$ bijectively and in an arbitrary order. In this way we ensure that f is a surjective homomorphism from G to H .

Now suppose that $f: V_G \rightarrow V_H$ is a surjective homomorphism. Because f is a homomorphism, f maps injectively every clique of G to a clique in H . Because $p_i \geq 3$ for all $1 \leq i \leq n$, we then find that f cannot map a clique U_i to an edge $x_q^{(j-1)}x_1^{(j)}$. Hence, f maps U_i injectively to some clique $X^{(j)}$ of H .

Let $1 \leq j \leq m$, and let $\{i_1, \dots, i_s\}$ be the set of all indices that correspond to the U -cliques that f maps to X_j . Suppose that $p_{i_1} + \dots + p_{i_s} < q$. Then, $6m^2(a_{i_1} + \dots + a_{i_s} + sB) = p_{i_1} + \dots + p_{i_s} < q = 24m^2B$. This means that $a_{i_1} + \dots + a_{i_s} + sB \leq 3$. Consequently, $q - (p_{i_1} + \dots + p_{i_s}) \geq 6m^2$. Hence, f maps at least $6m^2$ inner vertices of the paths Q_i to $X^{(j)}$. However, the total number of these vertices is $(n-1)(2m-2) = (3m-1)(2m-2) < 6m^2$, a contradiction. This means that $p_{i_1} + \dots + p_{i_s} \geq q$. Because the same claim holds for all $1 \leq j \leq m$, and $p_1 + \dots + p_n = 6m^2(a_1 + \dots + a_n + nB) = 6m^2(mB + 3mB) = 24m^3B = mq$, we conclude that $p_{i_1} + \dots + p_{i_s} = q$. Because $6m^2(a_{i_1} + \dots + a_{i_s} + sB) = p_{i_1} + \dots + p_{i_s} = q = 24m^2B$ and $a_{i_1} + \dots + a_{i_s} > 0$, we find that $s \leq 3$. Then, because the same claim holds for all $1 \leq j \leq m$, and $p_1 + \dots + p_n = 6m^2(a_1 + \dots + a_n + nB) = mq = 24m^3B$, we find that $s = 3$ and $a_{i_1} + a_{i_2} + a_{i_3} = B$. We set $S_j = \{a_{i_1}, a_{i_2}, a_{i_3}\}$. It remains to observe that S_1, \dots, S_m is a 3-partition of A . This completes the proof of (vii). \square

4 Tractable Cases

By Theorem 1 (v), SURJECTIVE HOMOMORPHISM is NP-complete when G and H are restricted to be trees. Here, we prove that the problem is FPT for trees when parameterized by the number of leaves in H . We first need some additional terminology. Let T be a tree. Then we may fix some vertex of T and call it the *root* of T . We observe that the root defines a parent-child relation between adjacent vertices. This enables us to define for a vertex u of T the tree G_u , which is the subtree of T that is induced by u and all its descendants in T ; we fix u to be the root of G_u . For a child v of u , we let G_{uv} denote the subtree of G induced by u and the set of all descendants of v in T ; we fix u to be the root of G_{uv} .

Theorem 2. *Testing if there is a surjective homomorphism from an n -vertex tree G to an m -vertex tree H with k leaves can be done in $O(2^{2k}nm^2)$ time.*




Figure 5: The graphs G and H as considered in the proof of Theorem 3.

Proof. We use dynamic programming. If H has one vertex the claim is trivial. Assume that H has at least one edge. Let L be the set of the leaves of H . First, we fix a root r of G . For each vertex $u \in V_G$, we construct a table that contains a number of records $R = (x, S)$ where $x \in V_H$ and $S \subseteq L$. A pair (x, S) is a record for u if and only if there exists a homomorphism h from G_u to H such that $h(u) = x$ and $S \subseteq h(V_{G_u})$. We also construct a similar table for each edge $uv \in E_G$. Then a pair (x, S) is a record for uv if and only if there exists a homomorphism h from G_{uv} to H such that $h(u) = x$ and $S \subseteq h(V_{G_{uv}})$. The key observation is that a homomorphism f from G to H is surjective if and only if $L \subseteq f(V_G)$, i.e., if and only if the table for r contains at least one record (z, L) .

We construct the tables as follows. We start with the leaves in G not equal to r (should r be a leaf). Their tables are constructed straightforwardly. Suppose that we have not constructed the table for a vertex u , while we have constructed the tables for all children v_1, \dots, v_p of u . Then we first determine the table for each edge uv_i by letting it consist of all records (x, S) such that

- (y, S) with $y \in N_H(x)$ is in the table for v_i ;
- $x \in L$ and $(y, S \setminus \{x\})$ with $y \in N_H(x)$ is in the table for v_i .

To construct the table for u , we consecutively construct auxiliary tables for $i = 1, \dots, p$. The table for $i = 1$ is the table for uv_1 . The table for $i \geq 2$ consists of the records (x, S) such that $S = S' \cup S''$, (x, S') is in the table for $i - 1$ and (x, S'') is in the table for uv_i . The table for u is the table constructed for $i = p$.

The correctness of the algorithm follows from its description. We observe that each table contains at most $m2^k$ records and can be constructed in $O(2^{2k} \cdot m^2)$ time. Because we construct $O(n)$ tables (including the auxiliary ones), our algorithm runs in $O(2^{2k} \cdot nm^2)$ time. This completes the proof of Theorem 2. \square \square

We now prove that SURJECTIVE HOMOMORPHISM is FPT when parameterized by the vertex cover number of G and H . The following approach has been successful before [8, 10]. The idea is to reduce a problem to an integer linear programming problem that is FPT when parameterized by the number of variables. Therefore, we consider the p -VARIABLE INTEGER LINEAR PROGRAMMING FEASIBILITY problem that has as input a $q \times p$ matrix A with integer elements and an integer vector $b \in \mathbb{Z}^q$ and that is to decide whether there exists a vector $x \in \mathbb{Z}^p$ such that $A \cdot x \leq b$. Lenstra [19] showed that this problem is FPT when parameterized by p . The best running time is due to Frank and Tardos [13].

Lemma 1 ([13]). *The p -VARIABLE INTEGER LINEAR PROGRAMMING FEASIBILITY problem can be solved using $O(p^{2.5p+o(p)} \cdot L)$ arithmetic operations and space polynomial in L , where L is the number of bits of the input.*

Theorem 3. *Testing if there is a surjective homomorphism from an n -vertex graph G with $\mathbf{vc}(G) \leq k$ to an m -vertex graph H with $\mathbf{vc}(H) \leq k$ can be done in $2^{2^{O(k)}}(nm)^{O(1)}$ time.*

Proof. Let G be an n -vertex graph with a vertex cover $S = \{u_1, \dots, u_s\}$ of size $s \leq k$. Then $I = V_G \setminus S$ is an independent set. For every subset $X \subseteq S$, we define $N(X)$ as the set of vertices in I that all have neighborhood X , i.e., $N(X) = \{u \in I \mid N(u) = X\}$. Note that $N(\emptyset)$ is the set of isolated vertices in I .

Let $X_1, \dots, X_r \subseteq S$ be the sets with $N(X_i) \neq \emptyset$. We let $p = s + r$ and define sets U_1, \dots, U_p where $U_i = \{u_i\}$ for $i = 1, \dots, s$ and $U_i = N(X_{i-s})$ for $i = s + 1, \dots, p$. We observe that $p \leq k + 2^k$ and that U_1, \dots, U_p is a partition of V_G , where each U_i is an independent set. Moreover, a vertex $v \in U_i$ is adjacent to a vertex $w \in U_j$ if and only if each vertex of U_i is adjacent to each vertex of U_j . In that case, we say that U_i is *adjacent* to U_j . We display G in Figure 5.

Let H be an m -vertex graph with a vertex cover $T = \{v_1, \dots, v_t\}$ of size $t \leq k$. Then $J = V_H \setminus T$ is an independent set, and for each $Y \subseteq T$ we define $N(Y) = \{z \in J \mid N(z) = Y\}$. Then we define $q \leq k + 2^k$ sets W_1, \dots, W_q where $W_j = \{v_j\}$ for $j = 1, \dots, t$ and $W_j = N(Y_{j-t})$ for $j = t + 1, \dots, q$. We also display H in Figure 5. The observations that we made for the U -sets are also valid for the W -sets.

Now we introduce integer variables x_{ij} for $1 \leq i \leq p$ and $1 \leq j \leq q$, and observe that there is a surjective mapping (not necessarily a homomorphism) $f: V_G \rightarrow V_H$ such that x_{ij} vertices of U_i are mapped to W_j if and only if the x_{ij} -variables satisfy the system

$$\begin{cases} x_{ij} \geq 0 & i \in \{1, \dots, p\}, j \in \{1, \dots, q\} \\ \sum_{j=1}^q x_{ij} = |U_i| & i \in \{1, \dots, p\} \\ \sum_{i=1}^p x_{ij} \geq |W_j| & j \in \{1, \dots, q\}. \end{cases} \quad (1)$$

The mapping f is a homomorphism from G to H if and only if the following holds: for each pair of variables $x_{ij}, x_{i'j'}$ such that $x_{ij} > 0$ and $x_{i'j'} > 0$, if U_i is adjacent to $U_{i'}$, then W_j is adjacent to $W_{j'}$.

We are now ready to give our algorithm. We first determine the set S and T . We then determine the U -sets and the W -sets. We guess a set R of indices (i, j) and only allow the variables x_{ij} for $(i, j) \in R$ to get non-zero value. Hence, we set $x_{ij} = 0$ for $(i, j) \notin R$. We then check whether for all pairs $(i, j), (i', j') \in R$, if U_i is adjacent to $U_{i'}$, then W_j is adjacent to $W_{j'}$. If not, then we discard R and guess a next one. Else we solve the system (1). If the system has an integer solution, then the algorithm returns YES; otherwise we try a next guess of R . If all guesses fail, then the algorithm returns No.

The correctness of the above algorithm follows from the aforementioned observations. We now estimate the running time. We can find S and T in time $1.2738^k n^{O(1)}$ and $1.2738^k m^{O(1)}$, respectively [3]. Then the sets U_1, \dots, U_p and W_1, \dots, W_q can be constructed in time $1.2738^k (nm)^{O(1)}$. The number of variables x_{ij} is $pq \leq (k + 2^k)^2 = 2^{O(k)}$. This means that there are at most $2^{2^{O(k)}}$ possibilities to choose R . By Lemma 1, system (1) (with some variables x_{ij} set to be zero) can be solved in time $2^{2^{O(k)}} (nm)^{O(1)}$. Hence, the total running time is $2^{2^{O(k)}} (nm)^{O(1)}$. This completes the proof of Theorem 3. \square \square

5 Conclusions

Our complexity study shows that the SURJECTIVE HOMOMORPHISM problem is already NP-complete on a number of very elementary graph classes such as linear forests, trees of small pathwidth, unions of complete graphs, cographs, split graphs and proper interval graphs. We conclude that there is not much hope for finding tractable results in this direction, and consider the computational complexity classification of the SURJECTIVE $(-, H)$ -HOMOMORPHISM problem as the main open problem; note that SURJECTIVE $(G, -)$ -HOMOMORPHISM is trivially polynomial-time solvable for any guest graph G .

As we observed in Section 1, the SURJECTIVE $(-, H)$ -HOMOMORPHISM problem is NP-complete already for any fixed host graph H that is nonbipartite. We also mentioned the existence of a bipartite graph H for which the problem is NP-complete [2] and that the problem can be solved in polynomial time whenever the host graph H is a fixed tree [14]. The paper of Feder et al. [7] on retractions provides a good starting point for the next step as we explain below.

A *pseudoforest* is a graph in which each connected component has at most one cycle. The RETRACTION problem is to test whether a graph G retracts to a graph H . Feder et al. [7] consider this problem for graphs

that may have self-loops. Applying their result to simple graphs yields the following. For any pseudoforest H , the $(-, H)$ -RETRACTION problem is NP-complete if H is nonbipartite or contains a cycle on at least 6 vertices, and it is polynomial-time solvable otherwise. It is an interesting open problem to show that $(-, H)$ -RETRACTION and SURJECTIVE $(-, H)$ -HOMOMORPHISM are polynomially equivalent for any fixed host graph H . All the evidence so far seems to suggest this.

References

- [1] A. Adiga, R. Chitnis, and S. Saurabh, Parameterized algorithms for boxicity, In: Proceedings of ISAAC 2010, LNCS 6506, 366–377, 2010.
- [2] M. Bodirsky, J. Kára and B. Martin, The complexity of surjective homomorphism problems – a survey, manuscript, ArXiv, <http://arxiv.org/abs/1104.5257>.
- [3] J. Chen, I. A. Kanj, and G. Xia, Improved Parameterized upper bounds for vertex cover, In: Proceedings of MFCS 2006, LNCS 4162, 238–249, 2006.
- [4] B. Courcelle and S. Olariu, Upper bounds to the clique width of graphs, Discrete Applied Mathematics 101 (2000), 77–114.
- [5] V. Dalmau, P.G. Kolaitis and M.Y. Vardi, Constraint satisfaction, bounded treewidth, and finite-variable logics, Proc. CP 2002, LNCS 2470, 223–254, 2006.
- [6] R. Enciso, M. R. Fellows, J. Guo, I. A. Kanj, F. A. Rosamond, and O. Suchý, What makes equitable connected partition easy, In: Proceedings of IWPEC 2009, LNCS 5917, 122–133, 2009.
- [7] T. Feder, P. Hell, P. Jonsson, A. Krokhin and G. Nordh, Retractions to pseudoforests, SIAM Journal on Discrete Mathematics 24 (2010), 101–112.
- [8] M. R. Fellows, D. Lokshtanov, N. Misra, F. A. Rosamond, and S. Saurabh, Graph layout problems parameterized by vertex cover, In: Proceedings of ISAAC 2008, LNCS 5369, 294–305, 2008.
- [9] J. Fiala and J. Kratochvíl, Locally constrained graph homomorphisms – structure, complexity, and applications, Computer Science Review 2 (2008), 97–111.
- [10] J. Fiala, P. A. Golovach, and J. Kratochvíl, Parameterized complexity of coloring problems: Treewidth versus vertex cover, Theor. Comp. Sci. 412 (2011), 2513–2523.
- [11] J. Fiala, and D. Paulusma, A complete complexity classification of the role assignment problem, Theor. Comp. Sci. 349 (2005), 67–81.
- [12] J. Flum and M. Grohe, Parameterized complexity theory, Texts in Theoretical Computer Science. An EATCS Series, Springer-Verlag, Berlin, 2006.
- [13] A. Frank and É. Tardos, An application of simultaneous Diophantine approximation in combinatorial optimization, Combinatorica 7 (1987), 49–65.
- [14] P. A. Golovach, D. Paulusma and J. Song, Computing vertex-surjective homomorphisms to partially reflexive trees, Proc. CSR 2011, LNCS 6651, 261–274, 2011.
- [15] M. R. Garey and D. R. Johnson, Computers and intractability, Freeman, NY, 1979.
- [16] M. Grohe, The complexity of homomorphism and constraint satisfaction problems seen from the other side, Journal of the ACM 54(2007).
- [17] P. Hell and J. Nešetřil, On the complexity of H -colouring, Journal of Combinatorial Theory, Series B 48 (1990), 92–110.

- [18] P. Hell and J. Nešetřil, *Graphs and homomorphisms*, Oxford University Pr., 2004.
- [19] H. W. Lenstra, Jr., Integer programming with a fixed number of variables, *Math. Oper. Res.* 8 (1983), 538–548.
- [20] B. Martin and D. Paulusma, The computational complexity of Disconnected Cut and 2K2-Partition, In: *Proceedings of CP 2011*, LNCS 6876, 561–575, 2011.
- [21] N. Vikas, Computational complexity of compaction to reflexive cycles, *SIAM Journal on Computing* 32, 253–280, 2002.
- [22] N. Vikas, Compaction, retraction, and constraint satisfaction, *SIAM Journal on Computing* 33 (2004), 761–782.
- [23] N. Vikas, A complete and equal computational complexity classification of compaction and retraction to all graphs with at most four vertices and some general results, *J. Comput. Syst. Sci.* 71 (2005), 406–439.
- [24] N. Vikas, Algorithms for partition of some class of graphs under compaction, In: *Proceedings of COCOON 2011*, LNCS 6842, 319–330, 2011.